

## Exercises 13      Applications of differential calculus Local/global maxima/minima, points of inflection

### Objectives

- be able to determine the local maxima and minima of a function.
- be able to determine the points of inflection of a function.
- be able to determine the global maximum and the global minimum of a cost, revenue, and profit function.
- be able to determine the global minimum of an average cost, average revenue, and average profit function.

### Problems

13.1 For each function, determine ...

- i) ... all local maxima and minima.
- ii) ... all points of inflection.
  
- a)  $f(x) = x^2 - 4$
- b)  $f(x) = -8x^3 + 12x^2 + 18x$
- c)  $s(t) = t^4 - 8t^2 + 16$
- d)  $f(x) = x e^{-x}$
- e) \*  $f(x) = (1 - e^{-2x})^2$
- f) \*  $V(r) = -D \left( \frac{2a}{r} - \frac{a^2}{r^2} \right)$       ( $D > 0, a > 0$ )

13.2 If the total profit for a commodity is

$$P(x) = (2000x + 20x^2 - x^3) \text{ CHF}$$

where  $x$  is the number of items sold, determine the level of sales,  $x$ , that maximises profit, and find the maximum profit.

Hints:

- First, find the local maxima.
- Then, check if one of the local maxima is the global maximum.

13.3 If the total cost for a service concerning a tourism event is given by

$$C(x) = \left( \frac{1}{4}x^2 + 4x + 100 \right) \cdot 100 \text{ CHF}$$

where  $x$  represents the extent of the service, what value of  $x$  will result in a minimum average cost? Determine the minimum average cost.

13.4 Suppose that the production capacity for a certain commodity cannot exceed 30. If the total profit for this company is

$$P(x) = (4x^3 - 210x^2 + 3600x) \text{ CHF}$$

where  $x$  is the number of units sold, determine the number of items that will maximise profit.

13.5 (see next page)

13.5 Suppose the annual profit for a store is given by

$$P(x) = (-0.1x^3 + 3x^2) \cdot 1000 \text{ CHF}$$

where  $x$  is the number of years past 2010. If this model is accurate, determine the point of inflection for the profit.

13.6 Decide which statements are true or false. Put a mark into the corresponding box. In each problem a) to c), exactly one statement is true.

a) If  $f$  has a local maximum at  $x = x_0$  it can be concluded that ...

- ...  $f(x_0) > f(x)$  for any  $x \neq x_0$
- ...  $f(x_0) > f(x)$  for any  $x > x_0$
- ...  $f(x_0) > f(x)$  for any  $x < x_0$
- ...  $f(x_0) > f(x)$  for all  $x$  which are in a certain neighbourhood of  $x_0$

b) If  $f(x_0) < 0$ ,  $f'(x_0) = 0$ , and  $f''(x_0) \neq 0$ , it can be concluded that  $f$  has ...


- ... no local minimum at  $x = x_0$
- ... no local maximum at  $x = x_0$
- ... no point of inflection at  $x = x_0$
- ... a point of inflection at  $x = x_0$

c) The global maximum of a function ...

- ... is always a local maximum.
- ... can be a local minimum.
- ... can be a local maximum.
- ... always exists.

**Answers**

13.1 a)  $f(x) = x^2 - 4$



$$f'(x) = 2x$$


$$f''(x) = 2$$

$$f'''(x) = 0$$

i)  $f'(x) = 0$  at  $x_1 = 0$   
 $f''(x_1) = 2 > 0$   $\Rightarrow$  local minimum at  $x_1 = 0$   
 no local maximum

ii)  $f''(x) = 2 \neq 0$  for all  $x$   $\Rightarrow$  no point of inflection

b)  $f(x) = -8x^3 + 12x^2 + 18x$



$$f'(x) = -24x^2 + 24x + 18$$


$$f''(x) = -48x + 24$$

$$f'''(x) = -48$$

i)  $f'(x) = 0$  at  $x_1 = -\frac{1}{2}$  and  $x_2 = \frac{3}{2}$   
 $f''(x_1) = 48 > 0$   $\Rightarrow$  local minimum at  $x_1 = -\frac{1}{2}$   
 $f''(x_2) = -48 < 0$   $\Rightarrow$  local maximum at  $x_2 = \frac{3}{2}$

ii)  $f''(x) = 0$  at  $x_3 = \frac{1}{2}$   
 $f'''(x_3) = -48 \neq 0 \quad \Rightarrow \quad \text{point of inflection at } x_3 = \frac{1}{2}$

c)  $s(t) = t^4 - 8t^2 + 16$




$s'(t) = 4t^3 - 16t$   
 $s''(t) = 12t^2 - 16$   
 $s'''(t) = 24t$

i)  $s'(t) = 0$  at  $t_1 = 0$ ,  $t_2 = -2$ , and  $t_3 = 2$   
 $s''(t_1) = -16 < 0 \quad \Rightarrow \quad \text{local maximum at } t_1 = 0$   
 $s''(t_2) = 32 > 0 \quad \Rightarrow \quad \text{local minimum at } t_2 = -2$   
 $s''(t_3) = 32 > 0 \quad \Rightarrow \quad \text{local minimum at } t_3 = 2$

ii)  $s''(t) = 0$  at  $t_4 = -\frac{2}{\sqrt{3}}$  and  $t_5 = \frac{2}{\sqrt{3}}$   
 $s'''(t_4) = -\frac{48}{\sqrt{3}} \neq 0 \quad \Rightarrow \quad \text{point of inflection at } t_4 = -\frac{2}{\sqrt{3}}$   
 $s'''(t_5) = \frac{48}{\sqrt{3}} \neq 0 \quad \Rightarrow \quad \text{point of inflection at } t_5 = \frac{2}{\sqrt{3}}$


d)  $f(x) = x e^{-x}$



$f'(x) = e^{-x} - x e^{-x} = (1 - x) e^{-x}$   
 $f''(x) = -e^{-x} - (1 - x) e^{-x} = (x - 2) e^{-x}$   
 $f'''(x) = e^{-x} - (x - 2) e^{-x} = (3 - x) e^{-x}$

- i)  $f'(x) = 0$  at  $x_1 = 1$   
 $f''(x_1) = -\frac{1}{e} < 0$   $\Rightarrow$  local maximum at  $x_1 = 1$   
no local minimum
- ii)  $f''(x) = 0$  at  $x_2 = 2$   
 $f'''(x_2) = \frac{1}{e^2} \neq 0$   $\Rightarrow$  point of inflection at  $x_2 = 2$

e) \*  $f(x) = (1 - e^{-2x})^2 = 1 - 2e^{-2x} + e^{-4x}$



$$f'(x) = 4e^{-2x} - 4e^{-4x} = 4e^{-2x}(1 - e^{-2x})$$

$$f''(x) = -8e^{-2x} + 16e^{-4x} = 8e^{-2x}(2e^{-2x} - 1)$$

$$f'''(x) = 16e^{-2x} - 64e^{-4x} = 16e^{-2x}(1 - 4e^{-2x})$$

- i)  $f'(x) = 0$  at  $x_1 = 0$   
 $f''(x_1) = 8 > 0$   $\Rightarrow$  local minimum at  $x_1 = 0$   
no local maximum
- ii)  $f''(x) = 0$  at  $x_2 = \frac{\ln(2)}{2} = 0.34\dots$   
 $f'''(x_2) = -8 \neq 0$   $\Rightarrow$  point of inflection at  $x_2 = 0.34\dots$

f) \*  $V'(r) = -D\left(-\frac{2a}{r^2} + \frac{2a^2}{r^3}\right) = \frac{2aD}{r^2}\left(1 - \frac{a}{r}\right)$   
 $V''(r) = -D\left(\frac{4a}{r^3} - \frac{6a^2}{r^4}\right) = \frac{2aD}{r^3}\left(\frac{3a}{r} - 2\right)$   
 $V'''(r) = -D\left(-\frac{12a}{r^4} + \frac{24a^2}{r^5}\right) = \frac{12aD}{r^4}\left(1 - \frac{2a}{r}\right)$

- i)  $V'(r) = 0$  at  $r_1 = a$   
 $V''(r_1) = \frac{2D}{a^2} > 0$   $\Rightarrow$  local minimum at  $r_1 = a$   
no local maximum
- ii)  $V''(r) = 0$  at  $r_2 = \frac{3a}{2}$   
 $V'''(r_2) = -\frac{64D}{81a^3} \neq 0$   $\Rightarrow$  point of inflection at  $r_2 = \frac{3a}{2}$

- 13.2 (Sole) local maximum at  $x_1 = \frac{100}{3} \rightarrow 33$  or 34  
 $P(33) = 51'843$  CHF  
 $P(34) = 51'816$  CHF  
 $P(x) < P(x_1)$  if  $x \neq x_1$  as there is no local minimum  
 $\Rightarrow P = 51'843$  CHF is the **global** maximum profit at  $x = 33$ .

13.3  $\bar{C}(x) = \frac{C(x)}{x} = \left(\frac{1}{4}x + 4 + \frac{100}{x}\right) \cdot 100 \text{ CHF}$

$\bar{C}(x)$  has a (sole) **local** minimum at  $x_1 = 20$ .

$\bar{C}(20) = 1400 \text{ CHF}$

$\bar{C}(x) > \bar{C}(x_1)$  if  $x \neq x_1$  as there is no local maximum.

$\Rightarrow \bar{C} = 1400 \text{ CHF}$  is the **global** minimum average cost at  $x = 20$ .

13.4  $P(x)$  has a **local** maximum at  $x_1 = 15$  and a **local** minimum at  $x_2 = 20$ .

$P(x_1) = 20'250 \text{ CHF}$

$P(x) < P(x_1)$  if  $x < x_1$  as there is no local minimum on the interval  $x < x_1$ .

$P(30) = 27'000 \text{ CHF} > 20'250 \text{ CHF} (!)$

$\Rightarrow P = 27'000 \text{ CHF}$  is the **global** maximum profit at the endpoint  $x = 30$ .

13.5  $P(x)$  has a point of inflection at  $x_1 = 10$ .

$P(10) = 200 \cdot 1000 \text{ CHF} = 200'000 \text{ CHF}$

$\Rightarrow$  point of inflection (10 | 200'000 CHF), i.e. when  $x = 10$  (in the year 2020) and  $P = 200'000 \text{ CHF}$

- 13.6
- a) 4<sup>th</sup> statement
  - b) 3<sup>rd</sup> statement
  - c) 3<sup>rd</sup> statement