
## **Derivative**

## **Function f**

 $f \colon \: D \to \mathbb{R} \qquad \quad \text{where } D \subseteq \mathbb{R}$ 

 $x \mapsto y = f(x)$ 

Ex.:  $f(x) = 24\sqrt{x+1} - 2x - 60$ 



What do we want to know?

**Slope of the tangent** to the graph of the function f at a certain point  $A(x_0 | f(x_0))$ .

Why do we want to know the slope?

- rate of change of values y with respect to x
- increasing (slope > 0), decreasing (slope < 0)
- local maximum/minimum (slope = 0)
- concavity (concave up if slope increases, concave down if slope decreases), points of inflection

Applications in economics


- tendency of costs/revenue/profit
- maximum/minimum of costs/revenue/profit
- marginal costs/revenue/profit: change of costs/revenue/profit if x (number of items or quantity of service) increases by one unit

## **Definition**

The slope of the tangent to the graph of f through the point  $A(x_0 | f(x_0))$  is called the **derivative** (or **rate of change**) **of f at x<sub>0</sub>**, denoted  $f'(x_0)$  ("f prime of  $x_0$ ").

**How** can we determine the slope?

The slope of the **secant** through the points  $A(x_0 | f(x_0))$  and  $B(x_0+\Delta x | f(x_0+\Delta x))$  tends towards the slope of the **tangent** through  $A(x_0 | f(x_0))$  as  $\Delta x$  tends towards 0.



Ex.: f: 
$$\mathbb{R} \to \mathbb{R}$$
  
 $x \mapsto y = f(x) = x^2$   
 $f'(x_0) = 2x_0$  (without proof)

Ex.: f: D \rightarrow \mathbb{R}  
 
$$x \mapsto y = f(x) = 24\sqrt{x+1} - 2x - 60$$
  
  $f'(x_0) = \frac{12}{\sqrt{x_0 + 1}} - 2$  (without proof)

## **Definition**

Suppose that the derivative (rate of change)  $f'(x_0)$  exists for all  $x_0 \in D_1$ , where  $D_1 \subseteq D$ .

The function f'

 $f': D_1 \to \mathbb{R}$ 

$$x \mapsto y = f'(x)$$

is called the derivative (or derived function) of f.

Ex.: f: 
$$\mathbb{R} \to \mathbb{R}$$
  
  $x \mapsto y = f(x) = x^2$ 

$$f': \mathbb{R} \to \mathbb{R}$$

$$x \mapsto y = f'(x) = 2x$$

Ex.: f: 
$$D \to \mathbb{R}$$

$$x \mapsto y = f(x) = 24\sqrt{x+1} - 2x - 60$$

$$f': D_1 \to \mathbb{R}$$

$$\begin{array}{ccc} f' \colon \ D_1 \to \mathbb{R} \\ & x \ \mapsto \ y = f'(x) = \frac{12}{\sqrt{x+1}} \ \text{-} \ 2 \end{array}$$

