

## Exercises 2                      **Function** **Domain, codomain, range, graph**


### Objectives

- understand what a function is.
- be able to judge whether a given relation is a function.
- be able to determine the range of a given function.
- be able to determine values of a given function.


### Problems

2.1 Which of the following relations are functions? Explain your answer.


a)




b)



c)



d)



- e) D = set of all courses in the FHGR Tourism bachelor programme  
 C = set of all FHGR lecturers  
 f:  $D \rightarrow C, c \mapsto l = f(c) = \text{lecturer of course } c$

- f)  $D = \{1994, 1995, \dots, 2003, 2004\}$   
 $C = \text{set of all human beings aged between 20 and 30}$   
 $f: D \rightarrow C, y \mapsto p = f(y) = \text{person who was born in the year } y$
- g)  $D = \text{set of all human beings aged between 20 and 30}$   
 $C = \{1994, 1995, \dots, 2003, 2004\}$   
 $f: D \rightarrow C, p \mapsto y = f(p) = \text{year of birth of person } p$
- h)  $f: \mathbb{R} \rightarrow \mathbb{R}, x \mapsto y = f(x) = x^2$
- i)  $f: \mathbb{R}^+ \rightarrow \mathbb{R}, x \mapsto y = f(x) = \text{number whose square is } x$
- Notice:  
 -  $\mathbb{R}^+$  is the set of all positive real numbers, i.e.  $\mathbb{R}^+ = \{x: x \in \mathbb{R} \text{ and } x > 0\}$ .
- j)  $f: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto b = f(t) = \text{bank account balance at time } t$

2.2 Determine the range  $R$  of the functions below:

- a)  $D = \{\text{January, February, March, ..., December}\}$   
 $C = \{A, B, C, \dots, Z\}$   
 $f: D \rightarrow C, m \mapsto l = f(m) = \text{initial letter of month } m$
- b)  $D = \text{set of all neighbouring countries of Switzerland}$   
 $C = \text{set of all European towns and cities}$   
 $c: D \rightarrow C, x \mapsto y = c(x) = \text{capital of neighbouring country } x$
- c) function  $f$  in problem 2.1 g)
- d) function  $f$  in problem 2.1 h)

2.3 a)  $f: \mathbb{R} \rightarrow \mathbb{R}, x \mapsto f(x) = x^3 - x$

Determine the following values:

- |              |               |                  |
|--------------|---------------|------------------|
| i) $f(1)$    | ii) $f(-2)$   | iii) $f(a)$      |
| iv) $f(b^2)$ | v) $f(a - b)$ | vi) $f(x^3 - x)$ |


b)  $g: \mathbb{R} \setminus \{-1\} \rightarrow \mathbb{R}, x \mapsto g(x) = \frac{x^2}{x+1}$

Determine the following values:

- |              |               |                                     |
|--------------|---------------|-------------------------------------|
| i) $g(2)$    | ii) $g(-3)$   | iii) $g(a)$                         |
| iv) $g(b^2)$ | v) $g(a - b)$ | vi) $g\left(\frac{x^2}{x+1}\right)$ |

2.4 (see next page)

2.4 The graph of a function  $f$  is given as follows:



- a) State the value of  $f(-1)$ .
- b) Estimate the value of  $f(2)$ .
- c) For what values of  $x$  is  $f(x) = 2$ ?
- d) Estimate the values of  $x$  such that  $f(x) = 0$ .
- e) State the domain  $D$  of  $f$ .
- f) State the range  $R$  of  $f$ .

2.5 Decide which statements are true or false. Put a mark into the corresponding box. In each problem a) to c), exactly one statement is true.

- a) The range of the function  $f: \{x: x \in \mathbb{R} \text{ and } x \geq 4\} \rightarrow \mathbb{R}, x \mapsto y = f(x) = \sqrt{x - 4}$ , is the set ...
  - ...  $\{x: x \in \mathbb{R} \text{ and } x \geq 4\}$
  - ...  $\{y: y \in \mathbb{R} \text{ and } y \geq 4\}$
  - ...  $\mathbb{R}$
  - ...  $\mathbb{R}_0^+$
- b)  $f$  cannot be a function if ...
  - ... the domain of  $f$  is no number set.
  - ... the codomain of  $f$  contains more elements than the domain of  $f$ .
  - ... the domain of  $f$  contains more elements than the codomain of  $f$ .
  - ... at least one element of the domain of  $f$  has more than one image.
- c) If the range of a function contains as many elements as the domain, it can be concluded that ...
  - ... the range is the same set as the domain.
  - ... the codomain contains as many elements as the domain.
  - ... each element of the codomain is also an element of the range.
  - ... no element of the range is associated to more than one element of the domain.

**Answers**

- 2.1 a) no function  
 No element (instead of exactly one element) of C is associated to one of the elements of D.
- b) function
- c) no function  
 Two elements (instead of exactly one element) of C are associated to one of the elements of D.
- d) function
- e) no function  
 More than one element (instead of exactly one element) of C are associated to some elements of D.
- f) no function  
 Many elements (instead of exactly one element) of C are associated to each element of D.
- g) function
- h) function
- i) no function  
 Two elements (instead of exactly one element) of  $\mathbb{R}$  are associated to each element of  $\mathbb{R}^+$ .
- j) function

- 2.2 a)  $R = \{A, D, F, J, M, N, O, S\}$
- b)  $R = \{\text{Berlin, Vienna, Vaduz, Rome, Paris}\}$
- c)  $R = C$
- d)  $R = \mathbb{R}_0^+$

Notice:

-  $\mathbb{R}_0^+$  is the set of all positive real numbers, including zero, i.e.  $\mathbb{R}_0^+ = \{x: x \in \mathbb{R} \text{ and } x \geq 0\}$ .

- 2.3 a) i)  $f(1) = 1^3 - 1 = 0$
- ii)  $f(-2) = (-2)^3 - (-2) = -6$
- iii)  $f(a) = a^3 - a$
- iv)  $f(b^2) = (b^2)^3 - b^2 = b^6 - b^2$
- v)  $f(a - b) = (a - b)^3 - (a - b) = a^3 - 3a^2b + 3ab^2 - b^3 - a + b$
- vi)  $f(x^3 - x) = (x^3 - x)^3 - (x^3 - x) = x^9 - 3x^7 + 3x^5 - 2x^3 + x$

- b) i)  $g(2) = \frac{2^2}{2+1} = \frac{4}{3}$
- ii)  $g(-3) = \frac{(-3)^2}{-3+1} = -\frac{9}{2}$
- iii)  $g(a) = \frac{a^2}{a+1}$
- iv)  $g(b^2) = \frac{(b^2)^2}{b^2+1} = \frac{b^4}{b^2+1}$
- v)  $g(a - b) = \frac{(a - b)^2}{(a - b) + 1} = \frac{a^2 - 2ab + b^2}{a - b + 1}$
- vi)  $g\left(\frac{x^2}{x+1}\right) = \frac{\left(\frac{x^2}{x+1}\right)^2}{\left(\frac{x^2}{x+1}\right) + 1} = \frac{x^4}{x^3 + 2x^2 + 2x + 1}$

- 2.4
- a)  $f(-1) = -2$
  - b)  $f(2) \approx 2.8$
  - c)  $x_1 = -3, x_2 = 1$
  - d)  $x_1 \approx -2.5, x_2 \approx 0.3$
  - e)  $D = \{x: x \in \mathbb{R} \text{ and } -3 \leq x \leq 3\}$
  - f)  $R = \{y: y \in \mathbb{R} \text{ and } -2 \leq y \leq 3\}$
- 2.5
- a) 4<sup>th</sup> statement
  - b) 4<sup>th</sup> statement
  - c) 4<sup>th</sup> statement