## **Exercises 11** Derivative


# Derivative (rate of change), derivative (derived function) of constant/power/exponential functions

### **Objectives**

- be able to estimate a derivative (rate of change) out of the graph of a function.
- be able to state the derivative (rate of change) of a constant and a linear function.
- be able to determine the derivative (derived function) of a constant and a linear function.
- be able to determine the derivative (derived function) of a basic power and a basic exponential function.
- be able to determine a derivative (rate of change) of a basic power and a basic exponential function.

#### **Problems**

11.1 The graph of a function f ist given as follows:



Estimate the derivative (rate of change)  $f'(x_0)$  at the given position  $x_0$ :

- a)  $x_0 = -1$
- b)  $x_0 = 0$
- c)  $x_0 = 1$
- d)  $x_0 = -2$

#### Hints:

- Draw the tangent to the graph of f at the given position  $x_0$ .
- Choose any two points on the tangent, and estimate their coordinates.
- Determine the slope of the tangent out of the estimated coordinates of the two points.

11.2 For each of the following functions f:  $\mathbb{R} \to \mathbb{R}$ ,  $x \mapsto y = f(x) = ...$ 

- i) ... draw the graph of f.
- ii) ... state the derivative (rate of change)  $f'(x_0)$  at the given position  $x_0$ .
- f(x) = 3a)

- $x_0 = 2$
- b)  $f(x) = c \ (c \in \mathbb{R})$
- any  $x_0 \in \mathbb{R}$

any  $x_0 \in \mathbb{R}$ 

f(x) = 2x - 3c)

- $x_0 = 4$
- d)  $f(x) = mx + q \ (m \in \mathbb{R} \setminus \{0\}, q \in \mathbb{R})$

Hint:

- If the graph of a function f is a straight line, the derivative (rate of change) f'(x<sub>0</sub>) is the slope of that straight line, i.e f'( $x_0$ ) has the same value at each position  $x_0$ , and therefore does not depend on  $x_0$ .

11.3 Determine f'(x):

- a)
- f(x) = 3
- f(x) = 0b)
- f(x) = -1c)

- $f(x) = x^3$ d)
- $f(x) = x^4$ e)
- $f(x) = x^5$ f)

- $f(\mathbf{x}) = \mathbf{x}^{17}$ g)
- h)  $f(x) = x^{200}$
- $f(x) = x^{100'001}$ i)

- $f(x) = x^{-1}$ i)
- $f(x) = x^{-2}$ k)
- $f(x) = x^{-17}$ 1)

- $f(x) = \frac{1}{x}$
- $f(x) = \frac{1}{x^3}$
- $f(x) = \frac{1}{x^{99}}$

- $f(x) = 3^x$ p)
- $f(x) = 5^x$ q)
- $f(x) = \left(\frac{2}{3}\right)^x$ r)

11.4 Determine the derivative (rate of change)  $f'(x_0)$  of the function f at the indicated position  $x_0$ :

- f(x) = x
  - i)  $x_0 = 0$
- ii)  $x_0 = 1$
- iii)  $x_0 = -2$

- $f(x) = x^5$ b)
  - $x_0 = 0$
- ii)
- iii)

- $f(x) = x^{-4}$ c)
  - $x_0 = -1$
- ii)
- iii)

- $f(x) = \left(\frac{2}{3}\right)^x$ d)
  - i)
- ii)  $x_0 = 1$
- iii)  $x_0 = -2$

11.5 Decide which statements are true or false. Put a mark into the corresponding box. In each problem a) to c), exactly one statement is true.

a) The derivative (rate of change) of a function f at the position  $x_0$  is a ...

> ... real number. ... function. ... tangent.

> > ... graph.

b) (see next page)

| b) | The derivative (derived function) f' of a function f is a |   |  |
|----|---|---|--|
|    |   | real number.  |  |
|    |   | function.   |  |
|    |   | tangent.  |  |
|    |   | graph.  |  |
| c) | f'(x <sub>0</sub> ) is                                    | $f'(x_0)$ is the slope of the   |  |
|    |   | secant through the points $(0 0)$ and $(x_0 f(x_0))$ .                          |  |
|    |   | secant through the points $(x_0+\Delta x f(x_0+\Delta x))$ and $(x_0 f(x_0))$ . |  |
|    |   | tangent to the graph of f through $(x_0 f(x_0))$ .                              |  |
|    |   | tangent to the graph of f' through $(x_0 f(x_0))$ .                             |  |