


# Quadratic function

## Definition

$f: D \rightarrow \mathbb{R}$	$(D \subseteq \mathbb{R})$
$x \mapsto y = f(x) = ax^2 + bx + c$ general form	$(a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}, c \in \mathbb{R})$
$y = f(x) = a(x - u)^2 + v$ vertex form	$(a \in \mathbb{R} \setminus \{0\}, u \in \mathbb{R}, v \in \mathbb{R})$


## Graph

1.  $y = f(x) = x^2$  ( $a = 1, u = 0, v = 0$ )



2. Parameter **u** (**u is varied**, a and v are kept constant)

$$\begin{aligned} y = f_0(x) &= x^2 & (a = 1, \mathbf{u} = \mathbf{0}, v = 0) \\ y = f_1(x) &= (x - 2)^2 & (a = 1, \mathbf{u} = \mathbf{2}, v = 0) \\ y = f_2(x) &= (x + 1)^2 & (a = 1, \mathbf{u} = \mathbf{-1}, v = 0) \end{aligned}$$




3. Parameter **v** (**v is varied**, a and u are kept constant)

$$y = f_0(x) = x^2 \quad (a = 1, u = 0, v = 0)$$

$$y = f_1(x) = x^2 + 3 \quad (a = 1, u = 0, v = 3)$$

$$y = f_2(x) = x^2 - 2 \quad (a = 1, u = 0, v = -2)$$




4. Parameter **a** (**a is varied**, u and v are kept constant)

$$y = f_0(x) = x^2 \quad (a = 1, u = 0, v = 0)$$


$$y = f_1(x) = 2x^2 \quad (a = 2, u = 0, v = 0)$$

$$y = f_2(x) = -2x^2 \quad (a = -2, u = 0, v = 0)$$



5. Parameter **a** (**a is varied**, u and v are kept constant)

$$\begin{aligned}
 y = f_0(x) &= x^2 & (\mathbf{a} = 1, u = 0, v = 0) \\
 y = f_1(x) &= \frac{1}{2}x^2 & (\mathbf{a} = \frac{1}{2}, u = 0, v = 0) \\
 y = f_2(x) &= -\frac{1}{2}x^2 & (\mathbf{a} = -\frac{1}{2}, u = 0, v = 0)
 \end{aligned}$$




6. The **graph** of a quadratic function is a **parabola**.

The parameter **a** determines the **shape** of the parabola, and whether the parabola opens upwards or downwards.

The parameters **u** and **v** determine the **position** of the parabola. They are the coordinates of the **vertex V** of the parabola:  $V(u|v)$

$$\begin{aligned}
 y = f_0(x) &= x^2 & (\mathbf{a} = 1, u = 0, v = 0) & & V(0|0) \\
 y = f_1(x) &= 2(x - 2)^2 - 4 & (\mathbf{a} = 2, u = 2, v = -4) & & V(2|-4) \\
 y = f_2(x) &= -\frac{1}{2}(x + 1)^2 + 6 & (\mathbf{a} = -\frac{1}{2}, u = -1, v = 6) & & V(-1|6)
 \end{aligned}$$




## Examples

1. Nature/Physics: Trajectories of water in a fountain



2. Geometry: Square




Area A at side length  $l$ :  $A = l^2$

$f: \mathbb{R}^+ \rightarrow \mathbb{R}$

$l \mapsto A = f(l) = l^2$       quadratic function

3. Physics: Free fall



Distance  $d$  after time  $t$ :  $d = \frac{1}{2}gt^2$       ( $g$  = gravity field strength)

$f: \mathbb{R} \rightarrow \mathbb{R}$

$t \mapsto d = f(t) = \frac{1}{2}gt^2$       quadratic function

4. Economics: Supply, Demand