


Exercises 3 **Function** **Domain, codomain, range, graph**

Objectives

- understand what a function is.
- be able to judge whether a given relation is a function.
- be able to determine the range of a given function.
- be able to determine values of a given function.

Problems

3.1 Which of the following relations are functions? Explain your answer.



- e) $D =$ set of all the modules of the HTW Tourism bachelor programme
 $B =$ set of all the HTW lecturers
 $f: D \rightarrow B, m \mapsto l = f(m) =$ lecturer of m
- f) $D = \{1986, 1987, \dots, 1995, 1996\}$
 $B =$ set of all the human beings aged between 20 and 30
 $f: D \rightarrow B, y \mapsto p = f(y) =$ person who was born in the year y
- g) $D =$ set of all the human beings aged between 20 and 30
 $B = \{1986, 1987, \dots, 1995, 1996\}$
 $f: D \rightarrow B, p \mapsto y = f(p) =$ year of birth of person p
- h) $f: \mathbb{R} \rightarrow \mathbb{R}, x \mapsto y = f(x) = x^2$
- i) $f: \mathbb{R}^+ \rightarrow \mathbb{R}, x \mapsto y = f(x) =$ number whose square is x
 Notice:
 - \mathbb{R}^+ is the set of all positive real numbers, i.e. $\mathbb{R}^+ = \{x: x \in \mathbb{R} \text{ and } x > 0\}$.
- j) $f: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto b = f(t) =$ bank account balance at time t

3.2 Determine the range E of the functions below:

- a) $D = \{\text{January, February, March, ..., December}\}$
 $B = \{A, B, C, ..., Z\}$
 $f: D \rightarrow B, m \mapsto l = f(m) = \text{initial letter of } m$
- b) $D = \text{set of all the neighbouring countries of Switzerland}$
 $B = \text{set of all the European cities}$
 $c: D \rightarrow B, x \mapsto y = c(x) = \text{capital of neighbouring country } x$
- c) function f in problem 3.1 g)
- d) function f in problem 3.1 h)

3.3 a) $f: \mathbb{R} \rightarrow \mathbb{R}, x \mapsto f(x) = x^3 - x$

Determine the following values:


- i) $f(1)$ ii) $f(-2)$ iii) $f(a)$
- iv) $f(b^2)$ v) $f(a - b)$ vi) $f(x^3 - x)$

b) $g: \mathbb{R} \setminus \{-1\} \rightarrow \mathbb{R}, x \mapsto g(x) = \frac{x^2}{x+1}$

Determine the following values:

- i) $g(2)$ ii) $g(-3)$ iii) $g(a)$
- iv) $g(b^2)$ v) $g(a - b)$ vi) $g\left(\frac{x^2}{x+1}\right)$

3.4 The graph of a function f is given as follows:



- a) State the value of $f(-1)$.
- b) Estimate the value of $f(2)$.
- c) For what values of x is $f(x) = 2$?
- d) Estimate the values of x such that $f(x) = 0$.
- e) State the domain D of f .
- f) State the range E of f .

3.5 Decide which statements are true or false. Put a mark into the corresponding box.
In each problem a) to c), exactly one statement is true.

a) The range of the function $f: \{x: x \in \mathbb{R} \text{ and } x \geq 4\} \rightarrow \mathbb{R}, x \mapsto y = f(x) = \sqrt{x-4}$, is the set ...

- ... $\{x: x \in \mathbb{R} \text{ and } x \geq 4\}$
- ... $\{y: y \in \mathbb{R} \text{ and } y \geq 4\}$
- ... \mathbb{R}
- ... \mathbb{R}_0^+

b) f cannot be a function if ...

- ... the domain of f is no number set.
- ... the codomain of f contains more elements than the domain of f .
- ... the domain of f contains more elements than the codomain of f .
- ... at least one element of the domain of f has more than one image.

c) If f is such that $f(x) = x$ for all elements x of the domain, it can be concluded that ...

- ... the domain of f is the same set as the codomain of f .
- ... the range of f is the same set as the codomain of f .
- ... the domain and the codomain of f contain the same number of elements.
- ... the domain and the range of f contain the same number of elements.

Answers

- 3.1 a) no function
 No element (instead of exactly one element) of B is assigned to one of the elements of D.
- b) function
- c) no function
 Two elements (instead of exactly one element) of B are assigned to one of the elements of D.
- d) function
- e) no function
 More than one element (instead of exactly one element) of B are assigned to some elements of D.
- f) no function
 Many elements (instead of exactly one element) of B are assigned to each element of D.
- g) function
- h) function
- i) no function
 Two elements (instead of exactly one element) of \mathbb{R} are assigned to each element of \mathbb{R}^+ .
- j) function

- 3.2 a) $E = \{A, D, F, J, M, N, O, S\}$
- b) $E = \{\text{Berlin, Vienna, Vaduz, Rome, Paris}\}$
- c) $E = B$
- d) $E = \mathbb{R}_0^+$

Notice:

- \mathbb{R}_0^+ is the set of all positive real numbers, including zero, i.e. $\mathbb{R}_0^+ = \{x: x \in \mathbb{R} \text{ and } x \geq 0\}$.

- 3.3 a) i) $f(1) = 1^3 - 1 = 0$
- ii) $f(-2) = (-2)^3 - (-2) = -6$
- iii) $f(a) = a^3 - a$
- iv) $f(b^2) = (b^2)^3 - b^2 = b^6 - b^2$
- v) $f(a - b) = (a - b)^3 - (a - b) = a^3 - 3a^2b + 3ab^2 - b^3 - a + b$
- vi) $f(x^3 - x) = (x^3 - x)^3 - (x^3 - x) = x^9 - 3x^7 + 3x^5 - 2x^3 + x$

- b) i) $g(2) = \frac{2^2}{2+1} = \frac{4}{3}$
- ii) $g(-3) = \frac{(-3)^2}{-3+1} = -\frac{9}{2}$
- iii) $g(a) = \frac{a^2}{a+1}$
- iv) $g(b^2) = \frac{(b^2)^2}{b^2+1} = \frac{b^4}{b^2+1}$
- v) $g(a - b) = \frac{(a - b)^2}{(a - b) + 1} = \frac{a^2 - 2ab + b^2}{a - b + 1}$
- vi) $g\left(\frac{x^2}{x+1}\right) = \frac{\left(\frac{x^2}{x+1}\right)^2}{\left(\frac{x^2}{x+1}\right) + 1} = \frac{x^4}{x^3 + 2x^2 + 2x + 1}$

- 3.4
- a) $f(-1) = -2$
 - b) $f(2) \approx 2.8$
 - c) $x_1 = -3, x_2 = 1$
 - d) $x_1 \approx -2.5, x_2 \approx 0.3$
 - e) $D = \{x: x \in \mathbb{R} \text{ and } -3 \leq x \leq 3\} = [-3, 3]$
 - f) $E = \{y: y \in \mathbb{R} \text{ and } -2 \leq y \leq 3\} = [-2, 3]$
- 3.5
- a) 4th statement
 - b) 4th statement
 - c) 4th statement