


## Exercises 13      Derivative Rate of change, derivative of constant/power/exponential functions

### Objectives

- be able to estimate a rate of change out of the graph of a function.
- be able to state the rate of change of a constant/linear function.
- be able to determine a rate of change of a basic power/exponential function.
- be able to determine the derivative of a constant/linear function.
- be able to determine the derivative of a basic power/exponential function.

### Problems

13.1    The graph of a function  $f$  is given as follows:



Estimate the rate of change  $f'(x_0)$  at the given position  $x_0$ :

- |    |            |    |            |
|----|------------|----|------------|
| a) | $x_0 = -1$ | b) | $x_0 = 0$  |
| c) | $x_0 = 1$  | d) | $x_0 = -2$ |

Hints:

- Draw the tangent to the graph of  $f$  at the given  $x_0$ .
- Estimate the slope of the tangent.

13.2 The graph of a constant or linear function is a straight line. Therefore, the “tangent” to the graph at any point of the graph is the graph itself.

For each of the following functions  $f: \mathbb{R} \rightarrow \mathbb{R}, x \mapsto y = f(x) = \dots$

- i) ... draw the graph of  $f$ .
- ii) ... state the rate of change  $f'(x_0)$  at the given  $x_0$ .
- a)  $f(x) = 3$   $x_0 = 2$
- b)  $f(x) = c$  ( $c \in \mathbb{R}$ ) any  $x_0 \in \mathbb{R}$
- c)  $f(x) = 2x - 3$   $x_0 = 4$
- d)  $f(x) = mx + q$  ( $m \in \mathbb{R} \setminus \{0\}, q \in \mathbb{R}$ ) any  $x_0 \in \mathbb{R}$
- e) \*  $f(x) = |x|$  any  $x_0 \in \mathbb{R}$

13.3 Determine  $f'(x)$ :

- |                         |                           |                              |
|-------------------------|---------------------------|------------------------------|
| a) $f(x) = 3$           | b) $f(x) = 0$             | c) $f(x) = -1$               |
| d) $f(x) = x^3$         | e) $f(x) = x^4$           | f) $f(x) = x^5$              |
| g) $f(x) = x^{17}$      | h) $f(x) = x^{200}$       | i) $f(x) = x^{100001}$       |
| j) $f(x) = x^{-1}$      | k) $f(x) = x^{-2}$        | l) $f(x) = x^{-17}$          |
| m) $f(x) = \frac{1}{x}$ | n) $f(x) = \frac{1}{x^3}$ | o) $f(x) = \frac{1}{x^{99}}$ |

13.4 Determine  $f'(x)$ :

- |  |  |  |
|--|--|--|
| a) $f(x) = 3^x$                        | b) $f(x) = 5^x$                          | c) $f(x) = 18^x$                         |
| d) $f(x) = \left(\frac{2}{3}\right)^x$ | e) $f(x) = \left(\frac{13}{17}\right)^x$ | f) $f(x) = \left(\frac{1}{4}\right)^x$   |
| g) $f(x) = \left(\frac{1}{e}\right)^x$ | h) * $f(x) = \left(\frac{3}{e}\right)^x$ | i) * $f(x) = \left(\frac{e}{3}\right)^x$ |

13.5 Determine the rate of change  $f'(x_0)$  of the function  $f$  at the indicated  $x_0$ :

- |  |                          |                           |
|--|--------------------------|---------------------------|
| a) $f(x) = x$                          | ii) $x_0 = 1$            | iii) $x_0 = -2$           |
| i) $x_0 = 0$                           |                          |                           |
| b) $f(x) = x^5$                        | ii) $x_0 = 2$            | iii) $x_0 = -\frac{2}{3}$ |
| i) $x_0 = 0$                           |                          |                           |
| c) $f(x) = x^{-4}$                     | ii) $x_0 = -\frac{4}{3}$ | iii) $x_0 = 0$            |
| i) $x_0 = -1$                          |                          |                           |
| d) $f(x) = \left(\frac{2}{3}\right)^x$ | ii) $x_0 = 1$            | iii) $x_0 = -2$           |
| i) $x_0 = 0$                           |                          |                           |

13.6 \* (see next page)

- 13.6 \* The rate of change  $f'(x_0)$  of  $f$  at  $x_0$  can be determined by looking at the secant through the points  $A(x_0 | f(x_0))$  and  $B(x_0 + \Delta x | f(x_0 + \Delta x))$  of the graph of  $f$ . The slope of this secant tends towards the slope of the tangent at  $A(x_0 | f(x_0))$  as  $\Delta x$  tends towards 0.

It has been shown in class how to determine  $f'(x_0)$  for the quadratic function  $f(x) = x^2$ .

Find  $f'(x_0)$  for the following functions  $f$ :

a)  $f(x) = x^3$                       b)  $f(x) = \frac{1}{x^2}$

- 13.7 Decide which statements are true or false. Put a mark into the corresponding box. In each problem a) to c), exactly one statement is true.

a) The rate of change of the function  $f$  at the position  $x_0$  is a ...

- ... real number.  
 ... function.  
 ... tangent.  
 ... graph.

b) The derivative  $f'$  of the function  $f$  is a ...

- ... real number.  
 ... function.  
 ... tangent.  
 ... graph.

c)  $f'(x_0)$  is the slope of the ...

- ... secant through the points  $(0|0)$  and  $(x_0|f(x_0))$ .  
 ... secant through the points  $(x_0 + \Delta x | f(x_0 + \Delta x))$  and  $(x_0 | f(x_0))$ .  
 ... tangent to the graph of  $f$  at  $(x_0 | f(x_0))$ .  
 ... tangent to the graph of  $f'$  at  $(x_0 | f(x_0))$ .

**Answers**

- 13.1 a)  $f'(-1) \approx 0$                       b)  $f'(0) \approx 2$   
 c)  $f'(1) \approx \frac{3}{2}$                               d)  $f'(-2) \approx -\frac{5}{3}$
- 13.2 a) i) ...  
           ii)  $f'(2) = 0$   
 b) i) ...  
           ii)  $f'(x_0) = 0$   
 c) i) ...  
           ii)  $f'(4) = 2$   
 d) i) ...  
           ii)  $f'(x_0) = m$   
 e) \* i) ...  
           ii)  $f'(x_0) = \begin{cases} 1 & (x_0 > 0) \\ -1 & (x_0 < 0) \\ \text{not defined} & (x_0 = 0) \end{cases}$
- 13.3 a)  $f'(x) = 0$                       b)  $f'(x) = 0$                       c)  $f'(x) = 0$   
 d)  $f'(x) = 3x^2$                       e)  $f'(x) = 4x^3$                       f)  $f'(x) = 5x^4$   
 g)  $f'(x) = 17x^{16}$                       h)  $f'(x) = 200x^{199}$                       i)  $f'(x) = 100'001x^{100'000}$   
 j)  $f'(x) = -x^{-2}$                       k)  $f'(x) = -2x^{-3}$                       l)  $f'(x) = -17x^{-18}$   
 m)  $f'(x) = -\frac{1}{x^2}$                       n)  $f'(x) = -\frac{3}{x^4}$                       o)  $f'(x) = -\frac{99}{x^{100}}$
- 13.4 a)  $f'(x) = 3^x \ln(3)$                       b)  $f'(x) = 5^x \ln(5)$                       c)  $f'(x) = 18^x \ln(18)$   
 d)  $f'(x) = \left(\frac{2}{3}\right)^x \ln\left(\frac{2}{3}\right)$                       e)  $f'(x) = \left(\frac{13}{17}\right)^x \ln\left(\frac{13}{17}\right)$   
 f)  $f'(x) = \left(\frac{1}{4}\right)^x \ln\left(\frac{1}{4}\right) = -\frac{\ln(4)}{4^x}$
- Hint:  
 - Logarithm rules (see formulary) can be applied in order to simplify the result.
- g)  $f'(x) = -\frac{1}{e^x}$                       h) \*  $f'(x) = \left(\frac{3}{e}\right)^x (\ln(3) - 1)$                       i) \*  $f'(x) = \left(\frac{e}{3}\right)^x (1 - \ln(3))$
- 13.5 a)  $f'(x) = 1$   
           i)  $f'(0) = 1$                       ii)  $f'(1) = 1$                       iii)  $f'(-2) = 1$   
 b)  $f'(x) = 5x^4$   
           i)  $f'(0) = 0$                       ii)  $f'(2) = 80$                       iii)  $f'\left(-\frac{2}{3}\right) = \frac{80}{81}$   
 c)  $f'(x) = -\frac{4}{x^5}$   
           i)  $f'(-1) = 4$                       ii)  $f'\left(-\frac{4}{3}\right) = \frac{243}{256}$                       iii)  $f'(0)$  is not defined  
 d) (see next page)

d)  $f'(x) = \left(\frac{2}{3}\right)^x \ln\left(\frac{2}{3}\right)$

i)  $f'(0) = \ln\left(\frac{2}{3}\right)$     ii)  $f'(1) = \frac{2}{3} \ln\left(\frac{2}{3}\right)$     iii)  $f'(-2) = \frac{9}{4} \ln\left(\frac{2}{3}\right)$

13.6 \* a)  $f'(x_0) = 3x_0^2$

b)  $f'(x_0) = -\frac{2}{x_0^3}$

13.7 a) 1<sup>st</sup> statement

b) 2<sup>nd</sup> statement

c) 3<sup>rd</sup> statement